

On WZ-pairs which prove Ramanujan series

Jesús Guillera

jguillera@gmail.com

Av. Cesáreo Alierta, 31 esc. izda 4^o-A, Zaragoza (Spain)

Abstract

The known WZ-proofs for Ramanujan-type series related to $1/\pi$ gave us the insight to develop a new proof strategy based on the WZ-method. Using this approach we are able to find more generalizations and discover first WZ-proofs for certain series of this type.

Keywords Ramanujan series for $1/\pi$. Hypergeometric series. WZ-method

2000 Mathematics Subject Classification Primary-33C20.

1 Introduction

The Ramanujan series for $1/\pi$ are of the form [3, pp. 352-354]

$$\sum_{n=0}^{\infty} z^n \frac{\left(\frac{1}{2}\right)_n (s)_n (1-s)_n}{(1)_n^3} (a + bn) = \frac{1}{\pi}, \quad (1)$$

where s determine the family according to the possible values $s = 1/2, 1/4, 1/3, 1/6$ and the parameters z, a, b are algebraic real numbers with $-1 \leq z < 1$. For an excellent survey on this kind of series, see [2], which in addition cites many works of the main contributors. The symbol $(x)_n$ which appears in the series is the rising factorial or Pochhammer symbol, defined for all $x \in \mathbb{C}$ by

$$(x)_n = \begin{cases} x(x+1) \cdots (x+n-1), & n \in \mathbb{Z}^+, \\ 1, & n = 0, \end{cases} \quad (2)$$

or more generally by

$$(x)_k = \frac{\Gamma(x+k)}{\Gamma(x)}. \quad (3)$$

For $k \in \mathbb{Z} - \mathbb{Z}^-$, (3) coincide with (2). But (3) is more general because it is also defined for all complex x and k such that $x + k \in \mathbb{C} - (\mathbb{Z} - \mathbb{Z}^+)$.

The Ramanujan series corresponding to rational values of the parameter z can be written in the form

$$\sum_{n=0}^{\infty} z^n \frac{\left(\frac{1}{2}\right)_n (s)_n (1-s)_n}{(1)_n^3} (a + bn) = \frac{c}{\pi}, \quad (4)$$

where a and b are integers and c^2 is a rational number. D. Zeilberger in [6] proved the series for $s = 1/2$, $z = -1$, $a = 1$, $b = 4$, $c = 2$ in a simple way, as an application of the WZ (Wilf and Zeilberger) method (Sect. 2). Motivated by this result, in [9] and [10] we used WZ-pairs to obtain WZ-demonstrable generalizations of the series in Table I. In this paper we give a new strategy for the WZ-method specifically designed to prove Ramanujan-type series.

s	z	a	b	c	s	z	a	b	c
1/2	-1	1	4	2	1/2	1/4	1	6	4
1/2	-1/8	1	6	$2\sqrt{2}$	1/2	1/64	5	42	16
1/4	-1/4	3	20	8	1/4	1/9	1	8	$2/\sqrt{3}$
1/4	-1/48	3	28	$16/\sqrt{3}$	1/6	-27/512	15	154	$32\sqrt{2}$

Table I

Using this strategy, we have found other WZ-demonstrable generalizations of the series in Table I and the first WZ-proofs of those in Table II. We recall that the first proofs of the Ramanujan-type series for $1/\pi$ were based on the theory of modular forms and modular equations. For example, the formulas in Table I corresponding to $z = 1/4$, $z = 1/64$, $z = -1/4$, $z = -1/48$ and $z = 1/9$ were discovered and proved in this way by S. Ramanujan [12, Eq.: 28, 29, 35, 36, 40]. See also [3, pp. 352-354].

s	z	a	b	c
1/3	1/2	1	6	$3\sqrt{3}$
1/3	-9/16	1	5	$4/\sqrt{3}$
1/3	-1/16	7	51	$12\sqrt{3}$

Table II

The first proofs of the identities in Table II were also based on the theory of modular forms, but were discovered and proved much later [5].

2 The WZ-method

We recall that a function $A(n, k)$ is hypergeometric in its two variables if the quotients

$$\frac{A(n+1, k)}{A(n, k)} \quad \text{and} \quad \frac{A(n, k+1)}{A(n, k)}$$

are rational functions in n and k , respectively. Also, a pair of hypergeometric functions $F(n, k)$ and $G(n, k)$ is said to be a Wilf and Zeilberger (WZ) pair [11, Chapt. 7] if

$$F(n+1, k) - F(n, k) = G(n, k+1) - G(n, k). \quad (5)$$

In this case, H. S. Wilf and D. Zeilberger [13] proved that there exists a rational function $C(n, k)$ such that

$$G(n, k) = C(n, k)F(n, k). \quad (6)$$

The rational function $C(n, k)$ is the, so-called, certificate of the pair (F, G) . To discover WZ-pairs, usually we use a Maple package written by D. Zeilberger, called EKHAD [11, Appendix A]. If it certifies a function, we have found a WZ-pair!. However in this paper we only look for WZ-pairs which follow a special pattern, and we do not need that package. If we sum (5) over all $n \geq 0$, we get

$$\sum_{n=0}^{\infty} G(n, k) - \sum_{n=0}^{\infty} G(n, k+1) = -F(0, k) + \lim_{n \rightarrow \infty} F(n, k). \quad (7)$$

If $\lim_{n \rightarrow \infty} F(n, k) = 0$ and $F(0, k) = 0$, we have the identity

$$\sum_{n=0}^{\infty} G(n, k) = \sum_{n=0}^{\infty} G(n, k+1).$$

If, in addition, the function

$$f(k) = \sum_{n=0}^{\infty} G(n, k) - C,$$

where C is a constant, satisfies the hypothesis of Carlson's theorem (see [1], p. 39), then we obtain

$$\sum_{n=0}^{\infty} G(n, k) = C, \quad \forall k \in \mathbb{C}.$$

Usually it is easy to determine the constant by choosing a particular value for k . In other cases, we can determine the constant by taking the limit as $k \rightarrow \infty$.

3 A new strategy

We give a strategy to prove Ramanujan-type series

$$\sum_{n=0}^{\infty} z^n B(n)(a + bn) = \frac{c}{\pi}, \quad (8)$$

with z , a and b rational.

Let $B(n, k)$ be a function such that $B(n, 0) = B(n)$ and hypergeometric in its two variables. We simplify the quotients $B(n+1, k)/B(n, k)$ and $B(n, k+1)/B(n, k)$ and denote the resulting denominators by $P(n, k)$ and $Q(n, k)$. Then we write

$$P(n, k) = P_r(n, k)P_{r'}(n, k), \quad Q(n, k) = Q_s(n, k)Q_{s'}(n, k),$$

where $P_r(n, k)$ and $Q_s(n, k)$ are the polynomials of greatest possible degrees r and s satisfying $P_r(-1, 0) \neq 0$ and $Q_s(0, -1) \neq 0$, respectively. Finally, we construct two rational functions $R(n, k)$ and $S(n, k)$ (of degree 1) in the following way:

$$R(n, k) = \frac{(a + bn)P_r(n, 0) + kU_r(n, k)}{P_r(n, k)}, \quad S(n, k) = \frac{nV_s(n, k)}{Q_s(n, k)}, \quad (9)$$

where

$$U_r(n, k) = \sum_{j=0}^r \sum_{i=0}^j d_{i,j-i} n^i k^{j-i}, \quad V_s(n, k) = \sum_{j=0}^s \sum_{i=0}^j e_{i,j-i} n^i k^{j-i}$$

are polynomials of degrees r and s whose coefficients must be determined. A careful observation of the WZ-pairs used to prove Ramanujan-type series in [9] and [10] shows that they are of the form

$$G(n, k) = z^n y^k B(n, k) R(n, k), \quad F(n, k) = z^n y^k B(n, k) S(n, k), \quad (10)$$

where $R(n, k)$ and $S(n, k)$ are rational functions of the form (9) that we can construct in the way explained. To discover more WZ-pairs of this type, we consider other examples of functions $B(n, k)$ and try to solve for y , d_{ij} , e_{ij} from the equation

$$G(n, k+1) - G(n, k) = F(n+1, k) - F(n, k),$$

that is, from

$$\frac{B(n, k+1)}{B(n, k)} R(n, k+1) y - R(n, k) = \frac{B(n+1, k)}{B(n, k)} S(n+1, k) z - S(n, k). \quad (11)$$

Simplifying (11), we obtain an equation of the form $H(n, k) = 0$, where H is a polynomial in n and k . If there are values of y , d_{ij} and e_{ij} such that all the coefficients of $H(n, k)$ are equal to zero, then by substituting them in $F(n, k)$ and $G(n, k)$ we get a WZ-pair. In addition, as $F(0, k) = 0$ we can apply the theory of the preceding section to prove that

$$\sum_{n=0}^{\infty} z^n y^k B(n, k) R(n, k) = \frac{c}{\pi}. \quad (12)$$

In that case, the function $B(n, k)$ is good enough to prove a Ramanujan series.

4 Types of searches

Let us define the functions (by symmetry, $j_2 = j_3$ when $s = 1/3, 1/4, 1/6$)

$$C(n, k) = \frac{\left(\frac{1}{2} + j_1 k\right)_n (s + j_2 k)_n ((1-s)_n + j_3 k)_n}{(1)_n (1 + j_4 k)_n (1_n + j_5 k)}, \quad D(k) = \frac{(t)_k (1-t)_k}{(1)_k^2}, \quad (13)$$

where t can be any of the numbers $1/2, 1/3, 1/4, 1/6$ and the j_i are suitable rational numbers. The simplest functions $B(n, k)$ that we have tried in this paper are of the form

$$B(n, k) = C(n, k) D(k). \quad (14)$$

In [9], we found some series involving the simpler function

$$D(k) = \frac{\left(\frac{1}{2}\right)_k}{(1)_k}, \quad (15)$$

and there are other candidates for $D(k)$ that we have not yet tried, for example

$$D(k) = \frac{\left(\frac{1}{4}\right)_k \left(\frac{3}{4}\right)_k \left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{\left(\frac{1}{2}\right)_k^2 (1)_k^2}. \quad (16)$$

When we could not find solutions of the form (14), we tried the longer expression

$$B(n, k) = C(n, k) \cdot \frac{\left(\frac{1}{2} + j_6 k\right)_n}{\left(\frac{1}{2} + j_7 k\right)_n} \cdot D(k), \quad (17)$$

Of course, we can take more factors and so there are many other possibilities. For example, we can consider

$$B(n, k) = C(n, k) \cdot \frac{\left(\frac{1}{4} + j_6 k\right)_n \left(\frac{3}{4} + j_6 k\right)_n}{\left(\frac{1}{4} + j_7 k\right)_n \left(\frac{3}{4} + j_7 k\right)_n} \cdot D(k). \quad (18)$$

However, we have not made the corresponding searches.

Our searches are not completely random, because there are functions that we can discard in advance. For example, the function

$$B(n, k) = \frac{\left(\frac{1}{2} - k\right)_n \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(1)_n (1+k)_n (1+2k)_n} \cdot \frac{\left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(1)_k^2} \quad (19)$$

cannot be good because it has a simple pole at $k = -1/2$, and so (12) cannot be finite at this value. Another example is the function

$$B(n, k) = \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2} + k\right)_n \left(\frac{1}{2} - 5k\right)_n}{(1)_n (1+k)_n^2} \cdot \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2}, \quad (20)$$

which cannot be good, because at $k = \frac{1}{10}$ all the summands of (12) are zero except those corresponding to $n = 0$ and we can easily check that we do not get a sum of the form c/π , where c^2 is a rational.

5 An example

Using our strategy, we prove in detail the following Ramanujan series in Table II:

$$\sum_{n=0}^{\infty} \left(\frac{-1}{16}\right)^n \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{(1)_n^3} (51n + 7) = \frac{12\sqrt{3}}{\pi}. \quad (21)$$

Solution 1: We will prove that the following function $B(n, k)$ is good:

$$B(n, k) = \frac{\left(\frac{1}{2} - k\right)_n \left(\frac{1}{2} + k\right)_n \left(\frac{1}{3}\right)_n \left(\frac{2}{3}\right)_n}{\left(\frac{1}{2} + \frac{k}{2}\right)_n \left(1 + \frac{k}{2}\right)_n (1+k)_n (1)_n} \cdot \frac{\left(\frac{1}{3}\right)_k \left(\frac{2}{3}\right)_k}{(1)_k^2}. \quad (22)$$

Proof: We define the functions

$$G(n, k) = \left(\frac{-1}{16}\right)^n y^k B(n, k) R(n, k), \quad F(n, k) = \left(\frac{-1}{16}\right)^n y^k B(n, k) S(n, k),$$

and get the evaluations

$$\frac{B(n+1, k)}{B(n, k)} = \frac{(2n-2k+1)(2n+2k+1)(3n+1)(3n+2)}{9(2n+k+1)(2n+k+2)(n+k+1)(n+1)}, \quad (23)$$

$$\frac{B(n, k+1)}{B(n, k)} = -\frac{(2n+2k+1)(3k+1)(3k+2)}{9(2n-2k-1)(2n+k+1)(n+k+1)}. \quad (24)$$

Applying our strategy, we write

$$R(n, k) = \frac{(51n + 7)(2n + 1) + k(d_{10}n + d_{01}k + d_{00})}{2n + k + 1}$$

and

$$S(n, k) = \frac{n(e_{10}n + e_{01}k + e_{00})}{2n - 2k - 1}.$$

Substituting in (11) and simplifying, we arrive at an equation of the form $H(n, k) = 0$, where H is a polynomial of degree 5 and so has 21 coefficients. As the coefficients of n^5 , n^4 , n^3 , n^2 , n^4k , nk^4 and of the independent term are, respectively,

$$\begin{aligned} & -58752 + 612e_{10}, \\ & -125568 + 1512e_{10} + 612e_{00}, \\ & -60192 + 1187e_{10} + 1476e_{00} - 6528y, \\ & 23328 + 278e_{10} + 1151e_{00} - 7424y - 64yd_{10}, \\ & -29376 + 1152e_{10} - 576d_{10} + 612e_{01}, \\ & -288yd_{10} + 288d_{10} + 144e_{01} - 288yd_{01} + 576d_{01}, \\ & 2016 - 2e_{10} - 224y - 2e_{00} - 32yd_{01} - 32yd_{00}, \end{aligned}$$

we immediately get $e_{10} = 96$, $e_{00} = -32$, $y = 1$, $d_{10} = 90$, $e_{01} = -48$, $d_{01} = 24$, $d_{00} = 28$. Substituting these values in the other coefficients, we see that all of them are zero and so the polynomial is identically zero. Thus, $B(n, k)$ is a good function and the solution is given by $y = 1$,

$$R(n, k) = \frac{(51n + 7)(2n + 1) + k(90n + 24k + 28)}{2n + k + 1}, \quad (25)$$

and

$$S(n, k) = \frac{16n(6n - 3k - 2)}{2n - 2k - 1}. \quad (26)$$

Then the WZ-method leads to

$$\sum_{n=0}^{\infty} G(n, k) = \sum_{n=0}^{\infty} G(n, k + 1),$$

and we can apply Carlson's theorem to derive

$$\sum_{n=0}^{\infty} G(n, k) = C, \quad \forall k \in \mathbb{C},$$

where we determine the constant C by taking $k = 1/2$. In this way, we have the formula

$$\sum_{n=0}^{\infty} \left(\frac{-1}{16} \right)^n \frac{\left(\frac{1}{2} - k \right)_n \left(\frac{1}{2} + k \right)_n \left(\frac{1}{3} \right)_n \left(\frac{2}{3} \right)_n}{\left(\frac{1}{2} + \frac{k}{2} \right)_n \left(1 + \frac{k}{2} \right)_n (1+k)_n (1)_n} \times \frac{(51n+7)(2n+1) + k(90n+24k+28)}{2n+k+1} = \frac{12\sqrt{3}}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{3} \right)_k \left(\frac{2}{3} \right)_k}. \quad (27)$$

Taking $k = 0$ we obtain (21).

Solution 2: Another good function $B(n, k)$ is

$$B(n, k) = \frac{\left(\frac{1}{2} \right)_n \left(\frac{1}{2} + 2k \right)_n \left(\frac{1}{3} + k \right)_n \left(\frac{2}{3} + k \right)_n}{\left(\frac{1}{2} + \frac{k}{2} \right)_n \left(1 + \frac{k}{2} \right)_n (1+k)_n (1)_n} \cdot \frac{\left(\frac{1}{4} \right)_k \left(\frac{3}{4} \right)_k}{(1)_k^2}. \quad (28)$$

Proof: We define the functions

$$G(n, k) = \left(\frac{-1}{16} \right)^n y^k B(n, k) R(n, k), \quad F(n, k) = \left(\frac{-1}{16} \right)^n y^k B(n, k) S(n, k).$$

Applying our strategy, we obtain $y = 1$,

$$R(n, k) = \frac{(51n+7)(2n+1) + k(114n+36k+37)}{2n+k+1}, \quad (29)$$

and

$$S(n, k) = \frac{-9n(6n^2 + 30nk + 13n - 7k - 3)}{(3k+1)(3k+2)}, \quad (30)$$

The WZ method leads to

$$\sum_{n=0}^{\infty} G(n, k) = \sum_{n=0}^{\infty} G(n, k+1),$$

and we can apply the Carlson's theorem to derive

$$\sum_{n=0}^{\infty} G(n, k) = C, \quad \forall k \in \mathbb{C},$$

where we determine the constant C by substituting $k = -1/3$. We can also find the constant by observing that as a consequence of the Weierstrass M-test [14, p. 49], the convergence of the series is uniform. Hence the following steps hold:

$$\lim_{k \rightarrow \infty} \sum_{n=0}^{\infty} G(n, k) = \sum_{n=0}^{\infty} \lim_{k \rightarrow \infty} G(n, k) = \frac{18\sqrt{2}}{\pi} \sum_{n=0}^{\infty} \left(\frac{-1}{8} \right)^n \binom{2n}{n} = \frac{12\sqrt{3}}{\pi},$$

where we have used the identity [4, p. 386]

$$\sum_{n=0}^{\infty} z^n \binom{2n}{n} = \frac{1}{\sqrt{1-4z}}.$$

Thus we have proved the formula

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\frac{-1}{16} \right)^n \frac{\left(\frac{1}{2} \right)_n \left(\frac{1}{2} + 2k \right)_n \left(\frac{1}{3} + k \right)_n \left(\frac{2}{3} + k \right)_n}{\left(\frac{1}{2} + \frac{k}{2} \right)_n (1 + \frac{k}{2})_n (1 + k)_n (1)_n} \\ \times \frac{(51n + 7)(2n + 1) + k(114n + 36k + 37)}{2n + k + 1} = \frac{12\sqrt{3}}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{4} \right)_k \left(\frac{3}{4} \right)_k}. \end{aligned} \quad (31)$$

Taking $k = 0$, we obtain (21).

6 More formulas

For each of the indicated Ramanujan series (see Tables I and II), we have chosen an example among the WZ-demonstrable generalizations that we have found using our method.

Case: $s = 1/2$, $z = -1$, $a = 1$, $b = 4$, $c = 2$.

$$\sum_{n=0}^{\infty} (-1)^n \frac{\left(\frac{1}{2} - k \right)_n^2 \left(\frac{1}{2} \right)_n (4n + 1)}{(1 + k)_n^2 (1)_n} = \frac{2}{\pi} \cdot \left(\frac{1}{4} \right)^k \cdot \frac{(1)_k^2}{\left(\frac{1}{4} \right)_k \left(\frac{3}{4} \right)_k}. \quad (32)$$

Case: $s = 1/2$, $z = 1/4$, $a = 1$, $b = 6$, $c = 4$.

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\frac{1}{4} \right)^n \frac{\left(\frac{1}{2} - k \right)_n \left(\frac{1}{2} + k \right)_n \left(\frac{1}{2} + 3k \right)_n (6n + 6k + 1)}{(1 + k)_n (1 + 2k)_n (1)_n} \\ = \frac{4}{\pi} \cdot \left(\frac{16}{27} \right)^k \cdot \frac{(1)_k^2}{\left(\frac{1}{6} \right)_k \left(\frac{5}{6} \right)_k}. \end{aligned} \quad (33)$$

Case: $s = 1/2$, $z = -1/8$, $a = 1$, $b = 6$, $c = 2\sqrt{2}$.

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\frac{-1}{8} \right)^n \frac{\left(\frac{1}{2} - k \right)_n \left(\frac{1}{2} + k \right)_n \left(\frac{1}{2} + 3k \right)_n (6n + 6k + 1)}{(1 + k)_n (1 + 2k)_n (1)_n} \\ = \frac{2\sqrt{2}}{\pi} \cdot \left(\frac{32}{27} \right)^k \cdot \frac{(1)_k^2}{\left(\frac{1}{6} \right)_k \left(\frac{5}{6} \right)_k}. \end{aligned} \quad (34)$$

Case: $s = 1/4$, $z = -1/4$, $a = 3$, $b = 20$, $c = 8$.

$$\sum_{n=0}^{\infty} \left(\frac{-1}{4} \right)^n \frac{\left(\frac{1}{2} + k \right)_n \left(\frac{1}{4} + \frac{3k}{2} \right)_n \left(\frac{3}{4} + \frac{3k}{2} \right)_n (20n + 18k + 3)}{(1+k)_n^2 (1)_n} = \frac{8}{\pi} \cdot \left(\frac{16}{27} \right)^k \cdot \frac{(1)_k^2}{\left(\frac{1}{6} \right)_k \left(\frac{5}{6} \right)_k}. \quad (35)$$

Case: $s = 1/4$, $z = 1/9$, $a = 1$, $b = 8$, $c = 2\sqrt{3}$.

$$\sum_{n=0}^{\infty} \left(\frac{1}{9} \right)^n \frac{\left(\frac{1}{2} \right)_n \left(\frac{1}{4} + \frac{3k}{2} \right)_n \left(\frac{3}{4} + \frac{3k}{2} \right)_n (8n + 6k + 1)}{(1+k)_n^2 (1)_n} = \frac{2\sqrt{3}}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{6} \right)_k \left(\frac{5}{6} \right)_k}. \quad (36)$$

Case: $s = 1/3$, $z = 1/2$, $a = 1$, $b = 6$, $c = 3\sqrt{3}$.

$$\sum_{n=0}^{\infty} \left(\frac{1}{2} \right)^n \frac{\left(\frac{1}{2} + k \right)_n \left(\frac{1}{3} \right)_n \left(\frac{2}{3} \right)_n (6n + 6k + 1)}{(1+k)_n (1+2k)_n (1)_n} = \frac{3\sqrt{3}}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{3} \right)_k \left(\frac{2}{3} \right)_k}. \quad (37)$$

Case: $s = 1/2$, $z = 1/64$, $a = 5$, $b = 42$, $c = 16$.

$$\sum_{n=0}^{\infty} \left(\frac{1}{64} \right)^n \frac{\left(\frac{1}{2} - k \right)_n \left(\frac{1}{2} \right)_n \left(\frac{1}{2} + k \right)_n \left(\frac{1}{2} + 2k \right)_n}{\left(\frac{1}{2} + \frac{k}{2} \right)_n \left(1 + \frac{k}{2} \right)_n (1+k)_n (1)_n} \times \frac{(42n + 5)(2n + 1) + k(84n + 24k + 26)}{2n + k + 1} = \frac{16}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{4} \right)_k \left(\frac{3}{4} \right)_k}. \quad (38)$$

Case: $s = 1/3$, $z = -1/16$, $a = 7$, $b = 51$, $c = 12\sqrt{3}$.

$$\sum_{n=0}^{\infty} \left(\frac{-1}{16} \right)^n \frac{\left(\frac{1}{2} \right)_n \left(\frac{1}{2} + 2k \right)_n \left(\frac{1}{3} + k \right)_n \left(\frac{2}{3} + k \right)_n}{\left(\frac{1}{2} + \frac{k}{2} \right)_n \left(1 + \frac{k}{2} \right)_n (1+k)_n (1)_n} \times \frac{(51n + 7)(2n + 1) + k(114n + 36k + 37)}{2n + k + 1} = \frac{12\sqrt{3}}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{4} \right)_k \left(\frac{3}{4} \right)_k}. \quad (39)$$

Case: $s = 1/3$, $z = -9/16$, $a = 1$, $b = 5$, $c = 4/\sqrt{3}$.

$$\sum_{n=0}^{\infty} \left(\frac{-9}{16} \right)^n \frac{\left(\frac{1}{2} - k \right)_n \left(\frac{1}{2} + 3k \right)_n \left(\frac{1}{3} + k \right)_n \left(\frac{2}{3} + k \right)_n}{\left(\frac{1}{2} \right)_n (1)_n (1+k)_n (1+3k)_n} \times \frac{(5n + 1)(2n + 1) + k(16n + 6k + 7)}{2n + 1} = \frac{4\sqrt{3}}{3\pi} \cdot 4^k \cdot \frac{(1)_k^2}{\left(\frac{1}{6} \right)_k \left(\frac{5}{6} \right)_k}. \quad (40)$$

Case: $s = 1/4$, $z = -1/48$, $a = 3$, $b = 28$, $c = 16\sqrt{3}$.

$$\sum_{n=0}^{\infty} \left(\frac{-1}{48} \right)^n \frac{\left(\frac{1}{2} - k\right)_n \left(\frac{1}{2} + 3k\right)_n \left(\frac{1}{4}\right)_n \left(\frac{3}{4}\right)_n}{\left(\frac{1}{2}\right)_n (1)_n (1+k)_n^2} \times \frac{(28n+3)(2n+1) + k(40n+18)}{2n+1} = \frac{16\sqrt{3}}{\pi} \cdot \frac{(1)_k^2}{\left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}. \quad (41)$$

Case: $s = 1/6$, $z = -27/512$, $a = 15$, $b = 154$, $c = 32\sqrt{2}$.

$$\sum_{n=0}^{\infty} \left(\frac{-27}{512} \right)^n \frac{\left(\frac{1}{2} - k\right)_n \left(\frac{1}{2} + k\right)_n \left(\frac{1}{6} + k\right)_n \left(\frac{5}{6} + k\right)_n}{\left(\frac{1}{2} + \frac{k}{2}\right)_n \left(1 + \frac{k}{2}\right)_n (1+k)_n (1)_n} \times \frac{(154n+15)(2n+1) + k(352n+108k+108)}{2n+k+1} = \frac{32\sqrt{2}}{\pi} \cdot \left(\frac{32}{27} \right)^k \cdot \frac{(1)_k^2}{\left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}. \quad (42)$$

We have checked that the WZ-pairs used in [7], [9] and [10] to prove similar series for $1/\pi^2$ satisfy the same criteria, except that the degree of the functions $R(n, k)$ and $S(n, k)$ is now 2. So we can try to use this strategy to prove the other unproved series in [8].

References

- [1] Bailey, W.N.: Generalized Hypergeometric Series. Cambridge Univ. Press, (1935).
- [2] Baruah, N.D., Berndt B.C., Chan, H.H.: Ramanujan's series for $1/\pi$: A survey, The Amer. Math. Monthly **116** (2009) 567-587.; also available at the pages <http://www.math.ilstu.edu/cve/speakers/Berndt-CVE-Talk.pdf>, and <http://www.math.uiuc.edu/berndt/articles/monthly567-587.pdf>.
- [3] Berndt, B.C.: Ramanujan's Notebooks, Part IV. Springer-Verlag, New York, (1994).
- [4] Borwein, J.M., Borwein, P.B.: Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity, (Canadian Mathematical Society Series of Monographs and Advanced Texts), John Wiley, New York, (1987).
- [5] Chan, H.H., Liaw, W.C., Tan, V.: Ramanujan's class invariant λ_n and a new class of series for $1/\pi$. J. London Math. Soc. **64**, 93-106, (2001).

- [6] Ekhad, S.B., Zeilberger D.: A WZ proof of Ramanujan's formula for π . In Raskias, J.M. (ed.). *Geometry, Analysis and Mechanics*. World Scientific, Singapore, 107-108, (1994); also available at arXiv:math/9306213v1. (The coauthor EKHAD is a Maple package written by D. Zeilberger).
- [7] Guillera, J.: Some binomial series obtained by the WZ-method. *Adv. in Appl. Math.* **29**, 599-603, (2002).
- [8] Guillera, J.: About a new kind of Ramanujan type series. *Exp. Math.* **12**, 507-510, (2003).
- [9] Guillera, J.: Generators of Some Ramanujan Formulas. *Ramanujan J.* **11**, 41-48, (2006).
- [10] Guillera, J.: *Series de Ramanujan: Generalizaciones y conjeturas*. Ph.D. Thesis, University of Zaragoza, Spain, (2007).
- [11] Petkovšek, M., Wilf, H.S., Zeilberger, D.: *A=B*. Peters A.K.: Ltd., (1996); also available at <http://www.math.upenn.edu/~wilf/AeqB.html>.
- [12] Ramanujan, S.: Modular equations and approximations to π . *Q. J. Math.* **45**, 350-372, (1914).
- [13] Wilf, H.S., Zeilberger, D.: Rational functions certify combinatorial identities. *Journal Amer. Math. Soc.* **3**, 147-158, (1990). (Winner of the Steele prize).
- [14] Whittaker, E.T., Watson, G.N.: *A Course of Modern Analysis*. Cambridge Univ. Press, (1927).